

THE TIRED BRAIN ON FIRE: THE RELATIONSHIP BETWEEN SLEEP DEPRIVATION, MIGRAINE, AND DEPRESSION IN YOUNG ADULTS

Dr. Anjali Raj , Medical Student, Samarkand state medical university, Uzbekistan

Pavithira Radhakrishnan , Medical Student, Samarkand state medical university, Uzbekistan

M. Bhagya Harini , Medical Student, Samarkand state medical university, Uzbekistan

Aftab saifi , Medical Student, Samarkand state medical university, Uzbekistan

Shariq Ali , Medical Student, Samarkand state medical university, Uzbekistan


Abstract


Sleep is often the first thing young adults sacrifice when academic demands, employment, social commitments, digital media, or irregular schedules become difficult to balance. Yet sleep is not simply a period of rest: it supports emotional regulation, pain processing, memory, and recovery. This narrative review examines the interconnected relationship between insufficient or disturbed sleep, migraine, and depressive symptoms in young adults. Available evidence suggests that the relationship is bidirectional. Poor sleep may lower the threshold for migraine attacks and is associated with greater headache burden, while migraine-related pain can further disturb sleep. Sleep disturbance is also closely linked with depression, and experimental evidence indicates that sleep loss can worsen emotional functioning. In college and university populations, poor sleep quality is common and is associated with depressive symptoms, although causality is difficult to establish from observational studies. Shared biological pathways involving hypothalamic and brainstem networks, circadian regulation, serotonin and other neuromodulatory systems, stress responses, and pain sensitivity may partly explain the overlap. The practical implication is not that every young adult with poor sleep will develop migraine or depression, but that persistent sleep problems deserve attention as part of a broader health assessment. Consistent sleep timing, adequate opportunity for sleep, management of migraine triggers, healthy daily routines, and timely professional support for persistent mood or headache symptoms may help reduce the burden.

Keywords: sleep deprivation; sleep quality; migraine; depression; depressive symptoms; young adults; insomnia; circadian rhythm; headache

1. Introduction

Young adulthood is a period of major change. Students may move away from home, begin demanding courses, work part-time or full-time, develop new relationships, and take responsibility





for their own routines. Sleep can become surprisingly easy to postpone. A late night followed by an early morning may seem manageable, but repeated sleep restriction can affect attention, mood, pain sensitivity, and daily functioning. A systematic review of college students found that poor sleep quality is common and that sleep quality is significantly associated with depression or depressive symptoms in this population [1].

Migraine adds another layer to the problem. People living with migraine frequently report that inadequate, fragmented, or irregular sleep can precede an attack, while the pain and associated symptoms of migraine can make normal sleep more difficult. Reviews describe this relationship as complex and bidirectional rather than a simple one-way pathway [2–4]. Depression is also more common among people with migraine than in the general population, and depressive symptoms may be associated with greater headache burden [5].


The phrase “tired brain on fire” is used here as a metaphor for the combination of sleep-related strain, heightened pain sensitivity, and emotional dysregulation. It should not be interpreted as literal brain inflammation in every person with sleep deprivation.


2. Sleep in Young Adulthood

Healthy sleep is shaped by both sleep pressure and the circadian clock. Young adults may naturally prefer later sleep and wake times, but early classes, work schedules, commuting, and social obligations can force a mismatch between biological timing and daily demands. This mismatch may result in shortened sleep or irregular sleep timing. Importantly, sleep duration is only one dimension of sleep health; quality, continuity, timing, and regularity also matter.

Sleep loss affects more than daytime tiredness. A 2024 systematic review and meta-analysis of experimental sleep-loss studies involving more than 5,700 participants found that sleep loss reduced positive affect and increased anxiety symptoms across studies, supporting the broader connection between inadequate sleep and emotional functioning [6]. This does not mean sleep deprivation alone causes clinical depression in every person, but it supports biological plausibility.

Sleep-related problem	Possible consequence	Relevance to migraine and mood
Short sleep duration	Daytime sleepiness, reduced attention, poorer emotional regulation	May reduce resilience to pain and stress.
Irregular sleep timing	Circadian disruption and inconsistent recovery	May act as a headache trigger for some people.





Fragmented sleep	Reduced restorative sleep and fatigue	Can coexist with migraine and mood symptoms.
Insomnia symptoms	Difficulty initiating or maintaining sleep	Closely linked with emotional distress and headache burden.
Large sleep-schedule shifts	Abrupt changes in routine	May trigger symptoms in susceptible individuals.

3. Sleep Deprivation and Migraine

Migraine is a neurological disorder characterized by recurrent attacks that may include moderate or severe headache, nausea, and sensitivity to light or sound. Sleep and migraine have an unusually close relationship. Reviews report that many people with migraine identify poor or insufficient sleep as a trigger, while sleep can also help terminate an attack [2–4].


The relationship is not simply “less sleep equals more migraine.” Both insufficient and excessive sleep, abrupt changes in routine, and poor sleep quality may be relevant for susceptible individuals. A person may sleep poorly because an attack is developing, then experience a migraine because disrupted sleep lowers their threshold for the next attack. This feedback loop can make it difficult to identify which problem came first.


Several mechanisms have been proposed. Brain regions involved in sleep–wake regulation overlap with networks implicated in migraine, while signaling systems involving serotonin, dopamine, melatonin, adenosine, and orexin may influence both sleep and headache biology. These mechanisms are still being studied and should be regarded as plausible contributors rather than a complete explanation [3,4].

4. Sleep Disturbance and Depression

The connection between sleep and depression is also bidirectional. Persistent sleep disturbance can precede the onset or recurrence of depression in some people, while depression can alter sleep timing, continuity, and perceived quality. Reviews describe sleep disturbance as more than a passive symptom in some cases; it may be a contributing risk factor and an important treatment target [7].

Evidence from young people is particularly relevant because delayed sleep–wake patterns and irregular schedules are common in this age group. A 2025 systematic review and meta-analysis of young individuals with delayed sleep–wake phase disorder found that most included studies reported greater depressive symptom severity among those with the disorder, although the evidence base was limited and varied in quality [8]. These findings support attention to circadian timing but do not establish that a late sleep schedule by itself causes depression.





It is important to distinguish depressive symptoms from a diagnosis of major depressive disorder. Feeling low, irritable, exhausted, or unmotivated after several nights of poor sleep does not automatically mean a person has depression. Persistent symptoms that interfere with school, work, relationships, or daily functioning warrant assessment by a qualified healthcare professional.

5. The Three-Way Relationship: Sleep, Migraine, and Depression

The overlap among these conditions is best understood as a network rather than three separate boxes. Poor sleep may increase pain sensitivity and emotional reactivity. Migraine can interrupt sleep through pain, nausea, or sensitivity to light and sound. Depression can alter motivation, daily structure, and sleep timing, while poor sleep can intensify emotional difficulties. Stress may influence all three at the same time.

A review of migraine, sleep, and emotional factors concluded that sleep disturbance and depression may both contribute to increased pain sensitivity and headache burden [2]. Research also suggests that depression and anxiety are important comorbidities in migraine and may be associated with more disabling headache [5].

Conceptual relationship among sleep, migraine and mood






Figure 1. Conceptual model of the bidirectional relationship among sleep disruption, migraine vulnerability, and depressive symptoms. This is a qualitative illustration, not measured effect sizes and not proof of causation.

6. Possible Biological Pathways

One proposed pathway involves altered pain processing. Sleep disruption can increase sensitivity to painful stimuli in some experimental and clinical contexts. In people already vulnerable to migraine, this may reduce the threshold at which an attack is triggered. Depression may further influence pain perception through emotional and central nervous system mechanisms [2].






A second pathway involves circadian and hypothalamic regulation. The hypothalamus participates in sleep–wake regulation and has also been implicated in migraine biology. Changes in melatonin and other circadian signals may therefore provide a bridge between irregular sleep and headache susceptibility [3,4].


A third pathway involves stress and emotional regulation. Repeated sleep restriction can make everyday stress feel harder to manage and can alter emotional responses. When combined with academic or social stress, poor sleep may contribute to a cycle of fatigue, reduced coping, and worsening symptoms.

Factor	Migraine relationship	Depression relationship	Interpretation
Insufficient sleep	May trigger or worsen attacks in susceptible people	Associated with poorer mood and emotional regulation	Assess pattern and persistence, not one bad night.
Poor sleep quality	Associated with greater headache burden	Strongly associated with depressive symptoms	Quality matters even when hours seem adequate.
Irregular schedule	Can be a trigger for some people	May disturb circadian regulation	Regular timing can be a useful behavioral target.
Migraine pain	Can delay or fragment sleep	Can increase emotional burden	Headache and sleep may need to be addressed together.
Depressive symptoms	Associated with greater migraine burden	May worsen sleep and motivation	Persistent symptoms need professional assessment.

7. Recognizing Warning Signs

A young adult should consider discussing symptoms with a healthcare professional when headaches become frequent or disabling, sleep problems persist despite reasonable routine changes, or low mood and loss of interest continue and interfere with daily life. Morning headaches accompanied by loud snoring, witnessed breathing pauses, or marked daytime sleepiness may also justify assessment for a sleep disorder.





Urgent medical assessment is appropriate for a sudden, unusually severe headache, a new neurological deficit, confusion, fainting, fever with a stiff neck, or a headache following significant head injury. These features should not be attributed to sleep deprivation without evaluation.


8. Practical Approaches


For young adults, the most useful approach is usually consistency rather than perfection. A regular wake time, adequate opportunity for sleep, morning exposure to natural light, daytime activity, and a predictable wind-down routine can support healthy sleep. If sleep problems continue, cognitive behavioral therapy for insomnia (CBT-I) is an evidence-based treatment that can be considered with professional guidance.

For migraine, a simple diary can help identify patterns such as irregular sleep, missed meals, dehydration, stress, or specific triggers. The purpose is not to create an increasingly restrictive lifestyle but to notice repeatable patterns and discuss them with a clinician. Migraine treatment should be individualized, particularly when headaches are frequent.

When depressive symptoms are present, young adults should not be expected to solve the problem through sleep hygiene alone. Support from a healthcare professional, counselor, or trusted adult may be appropriate. Sleep improvement can be part of care, but persistent depression requires a broader assessment and treatment plan.

Action	Practical idea	Goal
Protect sleep opportunity	Keep a fairly consistent sleep/wake schedule and allow enough time for sleep.	Reduce repeated sleep restriction.
Stabilize mornings	Use natural light and start the day at a consistent time when possible.	Support circadian regularity.
Track symptoms	Briefly record sleep timing, migraine days, and major mood changes.	Identify patterns without excessive monitoring.
Address persistent problems	Discuss ongoing insomnia, frequent migraine, or depressive symptoms with a professional.	Identify treatable conditions.
Avoid self-blame	Recognize that sleep problems can reflect	Use practical problem-solving instead of guilt.





	schedules, stress, illness, or environment.	
--	---	--

9. Discussion

The literature supports a meaningful association among sleep disturbance, migraine, and depression, but it should be interpreted carefully. Many studies are observational and cannot prove which condition causes another. People with migraine may sleep poorly because of pain, while people with depression may sleep poorly because of mood symptoms. Lifestyle factors such as stress, screen exposure, work schedules, caffeine use, physical activity, and socioeconomic circumstances can influence all three outcomes.


The strongest overall message is therefore not that sleep deprivation is a single cause of migraine or depression. Instead, sleep appears to be a modifiable part of a larger biological and behavioral system. Paying attention to sleep may help young adults and clinicians identify patterns earlier and may provide an accessible target for prevention and supportive care. Future research should prioritize longitudinal studies, objective sleep measurements, diverse young-adult populations, and interventions that examine headache and mood outcomes together.


10. Conclusion

The tired brain is not simply a metaphor for feeling sleepy. Repeated sleep disruption can influence emotional regulation, pain processing, attention, and daily functioning, while migraine and depression can in turn disrupt sleep. In young adults, the interaction may become a self-reinforcing cycle in which irregular sleep increases vulnerability to headache and emotional distress, while headache and low mood make restorative sleep harder to achieve. Current evidence supports a bidirectional relationship, but it does not justify assuming that sleep deprivation alone causes migraine or clinical depression. Consistent sleep routines, attention to migraine patterns, and early recognition of persistent mood or sleep problems are sensible components of a broader health strategy. When symptoms are severe, frequent, or persistent, professional assessment is important.

References

1. Dinis J, Bragança M. Quality of Sleep and Depression in College Students: A Systematic Review. *Sleep Sci.* 2018;11(4):290–301. doi:10.5935/1984-0063.20180045.
2. Rains JC, Poceta JS. Sleep disturbances and headache. *Headache.* 2006;46 Suppl 2:S68–S78.
3. Vgontzas A, Pavlović JM. Sleep Disorders and Migraine: Review of Literature and Potential Pathophysiology Mechanisms. *Headache.* 2018;58(7):1030–1039. doi:10.1111/head.13358.
4. Ferini-Strambi L, Galbiati A, Combi R. Sleep disorder-related headaches. *Neurol Sci.* 2014;35 Suppl 1:79–83.





5. Antonaci F, Nappi G, Galli F, et al. Migraine and psychiatric comorbidity: a review of clinical findings. *J Headache Pain*. 2011;12:115–125.
6. Palmer CA, et al. Sleep loss and emotion: A systematic review and meta-analysis of over 50 years of experimental research. *Psychol Bull*. 2024;150(3):306–335. doi:10.1037/bul0000410.
7. Alvaro PK, Roberts RM, Harris JK. A systematic review assessing bidirectionality between sleep disturbances and mental health problems. *Sleep Med Rev*. 2013;17(5):343–352.
8. The Association Between Delayed Sleep-Wake Phase Disorder and Depression Among Young Individuals: A systematic review and meta-analysis. *J Clin Sleep Med*. 2025.

