

VACCINE-PREVENTABLE DISEASE ELIMINATION

Purnima , Medical Student , Samarkand state medical university, Uzbekistan

Tanishqa, Medical Student , Samarkand state medical university, Uzbekistan

Khan Zainab , Medical Student , Samarkand state medical university, Uzbekistan

Ayuskha Mangesh Kadam , Medical Student , Samarkand state medical university, Uzbekistan


Abstract


Vaccine-preventable diseases have changed the history of public health. Diseases that once caused repeated childhood deaths, disability and fear can now be controlled, eliminated, and in the case of smallpox, eradicated through vaccination combined with strong surveillance and public-health systems. Yet elimination is not automatic. It depends on consistently high immunization coverage, equitable access, reliable disease surveillance, rapid outbreak response, public trust, strong supply chains and the ability of health systems to reach children who are missed by routine services. This narrative review examines the meaning of control, elimination and eradication; the role of routine immunization; lessons from polio, measles and rubella; barriers such as missed vaccinations, conflict, geographic access and vaccine confidence; and practical strategies for sustaining progress. Current WHO and UNICEF estimates show that global childhood vaccination has improved but remains below the level needed to achieve several elimination goals. In 2025, global DTP3 coverage was 85%, while approximately 13.5 million children received no routine vaccine dose. Measles-containing vaccine coverage also remained below the approximately 95% level generally needed to prevent sustained transmission. The review emphasizes that elimination is not simply a vaccine-delivery target: it is a long-term social and health-system achievement requiring every community to be reached and every outbreak to be detected quickly.

Keywords: vaccine-preventable diseases; immunization; disease elimination; eradication; measles; polio; rubella; vaccination coverage; surveillance; vaccine confidence; public health; primary healthcare.

1. Introduction

Vaccination is one of the clearest examples of prevention working before illness begins. Instead of waiting for a child to become seriously sick, immunization prepares the immune system in advance. Over decades, vaccination has reduced illness, disability and death from many infectious diseases





and has become a foundation of primary healthcare. WHO reports that immunization prevents millions of deaths every year and is central to global health security and outbreak prevention.


The phrase vaccine-preventable disease elimination can sound simple, but elimination is a demanding public-health goal. It means reducing transmission of a disease to zero in a defined geographic area through deliberate efforts, while continued measures may still be required to prevent re-establishment. Eradication is a still higher standard: permanent reduction of worldwide transmission to zero, after which routine interventions may no longer be needed. Smallpox remains the classic example of eradication. Polio is the major current eradication effort, while measles and rubella are prominent elimination goals.


The central lesson is that vaccines are necessary but not sufficient on their own. A country may have an effective vaccine but still experience outbreaks if children miss doses, communities are difficult to reach, surveillance is weak, or confidence in vaccination falls. Elimination therefore requires a complete system: vaccines, trained workers, cold-chain capacity, community engagement, laboratories, data, political commitment and rapid response.

2. Understanding Control, Elimination and Eradication

Term	Meaning	Example
Control	Reducing disease incidence or impact to an acceptable level through ongoing measures.	Long-term reduction of many vaccine-preventable infections through routine immunization.
Elimination	Reducing transmission or disease occurrence to zero in a defined geographic area, with continued measures needed to prevent reintroduction.	Measles or rubella elimination goals in specific regions.
Eradication	Permanent worldwide reduction of disease transmission to zero, making routine control measures unnecessary.	Smallpox eradication; global polio eradication remains an ongoing goal.

3. Why High Vaccination Coverage Matters





Population immunity is the bridge between individual vaccination and community protection. When a large proportion of people are immune, an infectious agent has fewer opportunities to move from one person to another. The exact coverage needed depends on how contagious the disease is and how well the vaccine protects against infection and transmission. Measles is particularly demanding because it spreads extremely efficiently; WHO's Immunization Agenda 2030 notes that around 95% coverage with two timely doses of measles-containing vaccine is needed to prevent sustained spread.

Current global figures show why elimination remains unfinished. WHO and UNICEF estimated that in 2025 about 90% of infants received at least one DTP-containing vaccine dose and 85% received the third dose. Approximately 13.5 million children were zero-dose children. UNICEF also reports that first-dose measles-containing vaccine coverage reached 86%, leaving millions of children without this protection. These are global averages, and averages can hide much lower coverage in particular communities or districts.






Figure 1. Selected global routine immunization coverage in 2025. Values are WHO/UNICEF estimates: DTP1 90%, DTP3 85%, and MCV1 86%.

4. The Role of Routine Immunization Systems

Elimination programs are strongest when they are built on reliable routine immunization rather than depending only on occasional campaigns. Routine services create repeated opportunities to vaccinate children at the recommended ages and to identify families who have missed appointments. They also





connect vaccination with other primary-healthcare services such as growth monitoring, nutrition support and health education.

A strong routine program requires more than vaccines arriving at a national warehouse. Doses must remain potent through the cold chain, health workers need appropriate training, vaccination sessions need to be convenient, records must be accurate, and communities need confidence that services are safe and respectful. WHO's Essential Programme on Immunization now supports vaccination across the life course, reflecting the fact that immunization is not only a childhood issue.

Equity is especially important. Children who are easiest to reach may already be well protected, while zero-dose and under-vaccinated children are often concentrated in remote, poor, displaced, conflict-affected or otherwise underserved communities. Elimination strategies must therefore ask not only 'What is the national coverage?' but also 'Who is still missing?'

5. Measles and Rubella: Lessons for Elimination


Measles is a useful indicator of the strength of an immunization system because its high transmissibility rapidly exposes immunity gaps. When coverage falls, outbreaks can occur even after years of progress. Rubella presents a different but equally important concern because infection during pregnancy can cause congenital rubella syndrome and severe birth defects. Preventing rubella transmission is therefore connected not only to childhood vaccination but also to protecting future pregnancies.


WHO reports that by January 2025, 179 of 194 countries had introduced rubella-containing vaccine, while global coverage was estimated at 73%. Although reported rubella cases have fallen dramatically since 2000, progress is uneven. For measles, regional elimination has also stalled in several settings. These experiences show that elimination cannot be declared and then forgotten. High coverage has to be sustained year after year, and imported infections must be detected and contained quickly.

A practical lesson is that elimination programs should treat every outbreak as information about the system. An outbreak can reveal a geographic pocket of low coverage, delayed vaccination, poor data quality, vaccine-access barriers or communication problems. Investigating these causes is more valuable than simply counting cases.

6. Polio and the Challenge of Eradication

Polio demonstrates both the power of vaccination and the difficulty of finishing an eradication effort. Decades of immunization have reduced poliovirus transmission enormously, but the final stage is demanding because any remaining transmission can threaten communities and allow spread across borders. WHO's 2025 mid-term review of Immunization Agenda 2030 noted that wild poliovirus





type 1 remained endemic in Afghanistan and Pakistan and that variant poliovirus transmission continued in several countries.

The polio experience highlights the importance of surveillance, laboratory capacity, rapid vaccination responses and community trust. Children who are repeatedly missed by vaccination teams are not simply statistics; they often represent practical barriers such as mobility, insecurity, inaccessible locations, misinformation, or gaps between health services and communities. Successful eradication therefore requires social and operational solutions as well as biological tools.

7. Barriers to Vaccine-Preventable Disease Elimination

The barriers to elimination are rarely caused by one factor. Access problems can occur when health facilities are distant, transport is difficult, vaccination sessions are irregular, or families cannot easily return for follow-up doses. Conflict and displacement can interrupt services and destroy health infrastructure. Outbreaks can also follow the disruption of routine programs after emergencies.


Vaccine confidence is another important component. People may have questions about safety, side effects, effectiveness or the need for vaccination when a disease becomes less visible. The most effective response is not ridicule or pressure. Health workers should listen to concerns, communicate clearly, explain benefits and risks honestly, and use trusted local voices. Confidence grows when people experience respectful care and consistent information.


Health-system barriers can be equally important. Stock-outs, inadequate cold-chain equipment, staff shortages, weak data systems and poor follow-up of missed children can reduce coverage even when communities are willing to vaccinate. This is why elimination is fundamentally a health-system strengthening challenge.

8. Surveillance and Outbreak Response

Vaccination coverage tells us how many people have received a vaccine; surveillance tells us what is happening with disease. Elimination requires both. Sensitive surveillance should identify suspected cases quickly, collect appropriate specimens, confirm diagnoses through reliable laboratories, map transmission and guide response.

Rapid response matters because vaccine-preventable infections can spread before a health system realizes that an outbreak has begun. Once an outbreak is detected, response may include case investigation, contact and community assessment, targeted vaccination, risk communication and strengthened routine services. The exact response depends on the disease, local epidemiology and national guidance.





Digital records and data systems can improve identification of children who miss doses, but technology is only useful when information is accurate and frontline workers can act on it. Data should be used to find gaps and improve services rather than simply to produce reports.

9. Community Engagement and Human Factors


Elimination is ultimately a community achievement. Health workers may deliver vaccines, but families decide whether and when to attend, community leaders influence trust, and local organizations can help identify children who are being missed. Programs that understand local routines, languages and concerns are more likely to reach people effectively.


A humanized approach recognizes that vaccine decisions are often shaped by real experiences. Parents may be balancing work, transport, childcare, previous experiences with healthcare and information from family or social networks. Listening before advising can make communication more effective. Respectful conversations also reduce the risk that people feel judged when they have questions.

The same principle applies to health workers. Staff need manageable workloads, adequate supplies, training and supportive supervision. If an immunization program expects excellent performance but does not provide the tools needed to achieve it, the gap is a system failure rather than simply an individual failure.

10. Strategies to Accelerate Elimination

Priority	Practical strategy	Expected benefit
Reach zero-dose children	Use local data to identify underserved settlements and provide convenient outreach or mobile services.	Improves equity and closes immunity gaps.
Strengthen routine services	Maintain vaccine supply, cold chain, trained staff and reliable session schedules.	Makes protection sustainable between campaigns.
Improve surveillance	Link community detection, laboratories and rapid reporting.	Finds outbreaks earlier and supports targeted action.
Build vaccine confidence	Listen to concerns and communicate with clear,	Supports informed acceptance and trust.





	respectful, evidence-based messages.	
Use outbreak response strategically	Investigate cases, map transmission and vaccinate according to national guidance.	Limits spread and reveals weaknesses.
Maintain political commitment	Fund immunization, surveillance and workforce capacity over the long term.	Prevents short-term gains from being lost.

Figure 2. Conceptual pathway from strong immunization systems to disease elimination

High routine coverage High population immunity Fewer outbreaks Interrupted transmission Elimination sustained





Figure 2. Conceptual pathway showing how sustained high coverage, population immunity, surveillance and rapid response support elimination. This is a conceptual model, not quantitative data.

11. Elimination, Equity and Primary Healthcare

Elimination programs can strengthen primary healthcare when they are designed to leave useful systems behind. Cold-chain improvements, trained vaccinators, disease surveillance, laboratory networks and better community outreach can support many other health programs. Conversely, an immunization campaign that operates completely separately from routine services may produce short-term coverage gains without fixing the underlying access problem.

Equity should therefore be treated as an outcome, not just a principle. National averages can improve while specific communities remain vulnerable. Disaggregated data by geography and other relevant factors can help programs locate these gaps. The goal is not merely to achieve a high national percentage; it is to ensure that protection reaches people who are most likely to be missed.





12. Discussion

The evidence and experience of global immunization programs point to a consistent conclusion: elimination is possible, but it is fragile. WHO's mid-term review of Immunization Agenda 2030 reported that the world was off track for several elimination and eradication goals and that large disruptive outbreaks of vaccine-preventable diseases remained frequent in 2024. At the same time, the continued expansion of vaccination programs and the large number of deaths prevented each year demonstrate that progress is achievable.

The most important shift is to view elimination as continuous prevention rather than a one-time finish line. High coverage must be maintained even when disease becomes rare. Surveillance must continue even when no cases are detected. Communities must remain engaged even after an outbreak ends. Health systems must be prepared to identify imported infections and respond before local transmission becomes established.

This perspective also makes the topic more human. Behind every percentage are children, parents, health workers and communities. A missed vaccine can mean an avoidable illness; a timely vaccine can mean that illness never happens. Elimination is therefore not only an epidemiological ambition but a commitment to protect people who may never know that prevention worked.

13. Conclusion


Vaccine-preventable disease elimination is one of public health's most ambitious and worthwhile goals. Vaccines provide the biological foundation, but elimination depends on much more: equitable access, strong routine immunization, reliable cold chains, trained health workers, accurate records, sensitive surveillance, laboratory confirmation, rapid outbreak response and sustained public trust.


Current global data show both progress and unfinished work. In 2025, global DTP3 coverage reached 85%, yet millions of children remained unvaccinated or incompletely vaccinated, and measles coverage remained below the level needed to reliably prevent transmission. These gaps remind us that elimination cannot be achieved by national averages alone. The hardest-to-reach communities must be reached deliberately and respectfully.

Ultimately, successful elimination is built through consistency. Vaccinate every eligible child, identify those who are missed, listen to communities, investigate every suspected outbreak, strengthen routine services and keep political commitment strong. When these actions continue for long enough, diseases that once seemed inevitable can become rare, then disappear from communities, and potentially disappear from the world.

References

1. World Health Organization. Vaccines and immunization. Geneva: WHO; 2026.





2. World Health Organization. Essential Programme on Immunization. Geneva: WHO; 2026.
3. World Health Organization. Immunization Agenda 2030: A Global Strategy to Leave No One Behind. Geneva: WHO; 2020.
4. World Health Organization. Immunization Agenda 2030: Mid-term review. Geneva: WHO; 2025.
5. World Health Organization. Routine vaccination coverage – worldwide, 2024. Weekly Epidemiological Record. 2025;100:627-641.
6. World Health Organization. WHO recommendations for routine immunization – summary tables. Geneva: WHO; updated December 2025.
7. World Health Organization. Rubella. Fact sheet. Geneva: WHO; 15 July 2026.
8. UNICEF. Vaccination and Immunization Statistics. UNICEF Data; 2026.
9. UNICEF and WHO. Global childhood immunization coverage inches forward despite conflict and hesitancy. 15 July 2026.
10. UNICEF and WHO. Measles cases dropped in Europe and Central Asia in 2025 compared to the previous year, but the risk of outbreaks remains. 2026.
11. World Health Organization. Smallpox and its eradication: history and lessons for global health. Geneva: WHO.
12. Hinman AR, Orenstein WA, Williamson DE, Darrington D. Childhood immunization: where we stand today. Vaccine. 2011;29 Suppl 1:C2-C6.

